

## Medline Abstract



Y Ueda and N Shimizu  
**Effects of pulse frequency of low-level laser therapy (LLLT) on bone nodule formation in rat calvarial cells.**  
J Clin Laser Med Surg, October 1, 2003; 21(5): 271-7.

- ▶ [Abstract](#)
- ▶ [Full text](#)
- ▶ [Order Full text via Infotrieve](#)
- ▶ [Alert Me when Cited](#)
- ▶ [MatchMaker](#)

- ▶ [Download to Citation Manager](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [PubMed Citation](#)
- ▶ [Related Articles in PubMed](#)
- ▶ [Order Full text via Infotrieve](#)
- ▶ [Publisher's Full Text Link](#)

Department of Orthodontics, Nihon University School of Dentistry at Matsudo Chiba, Japan.

**OBJECTIVE:** The purpose of this study was to determine the effect of pulse frequencies of low-level laser therapy (LLLT) on bone nodule formation in rat calvarial cells in vitro.  
**BACKGROUND DATA:** Various photo-biostimulatory effects of LLLT, including bone formation, were affected by some irradiation factors such as total energy dose, irradiation phase, laser spectrum, and power density. However, the effects of pulse frequencies used during laser irradiation on bone formation have not been elucidated. **MATERIALS AND METHODS:** Osteoblast-like cells isolated from fetal rat calvariae were irradiated once with a low-energy Ga-Al-As laser (830 nm, 500 mW, 0.48-3.84 J/cm<sup>2</sup>) in four different irradiation modes: continuous irradiation (CI), and 1-, 2-, and 8-Hz pulsed irradiation (PI-1, PI-2, PI-8). We then investigated the effects on cellular proliferation, bone nodule formation, alkaline phosphatase (ALP) activity, and ALP gene expression. **RESULTS:** Laser irradiation in all four groups significantly stimulated cellular proliferation, bone nodule formation, ALP activity, and ALP gene expression, as compared with the non-irradiation group. Notably, PI-1 and -2 irradiation markedly stimulated these factors, when compared with the CI and PI-8 groups, and PI-2 irradiation was the best approach for bone nodule formation in the present experimental conditions. **CONCLUSION:** Since low-frequency pulsed laser irradiation significantly stimulates bone formation in vitro, it is most likely that the pulse frequency of LLLT an important factor affecting biological responses in bone formation.

Publication Type:

- Journal article

MeSH Terms:

- Animals
- Cell Culture
- Fetus
- Laser Therapy, Low-Level
- Osteoblasts
- Osteogenesis
- Rats
- Skull
- Support, Non-U.S. Gov't

PMID: 14651794

MEDLINE data is licensed by HighWire Press from the National Library of Medicine. Some material in the NLM databases is from copyrighted publications of the respective copyright claimants. Users of the NLM databases are solely responsible for compliance with any copyright restrictions and are referred to the publication data appearing in the bibliographic citations, as well as to the copyright notices appearing in the original publications, all of which are hereby incorporated by reference.